

Reflected BSDEs with general filtration and two completely separated barriers

Mateusz Topolewski

Faculty of mathematics and Computer Science, Nicolaus Copernicus University,
Chopina 12/18, 87–100 Toruń, Poland.
E-mail: woland@mat.umk.pl

Abstract

We consider reflected backward stochastic differential equations, with two barriers, defined on probability spaces equipped with filtration satisfying only the usual assumptions of right continuity and completeness. As for barriers we assume that there are càdlàg processes of class D that are completely separated. We prove the existence and uniqueness of solutions for integrable final condition and integrable monotone generator. An application to zero-sum Dynkin game is given.

MSC: Primary 60H10; Secondary 60H30, 60H99.

Keywords: Reflected BSDE; general filtration; completely separated barriers; L^1 data

1 Introduction and notation

In this paper we study the problem of existence and uniqueness of solutions of backward stochastic differential equations with two reflecting càdlàg barriers L, U . The main new feature is that we deal with equations on probability spaces with general filtration $\mathbb{F} = \{\mathcal{F}_t; t \in [0, T]\}$ satisfying only the usual conditions of right-continuity and completeness, and we do not assume that the barriers satisfy so-called Mokobodzki condition. Instead, we assume that the lower barrier L and the upper barrier U are completely separated in the sense that $U_t < L_t$ and $U_{t-} < L_{t-}$ for $t \in [0, T]$. Moreover, we consider equations with L^p data, where $p \in [1, 2]$. Our motivation for considering such general setting comes from PDEs theory (equations involving nonlocal operators, see [9, 11]) and from the theory of optimal stopping (Dynkin games, see [8, 12, 14, 15]).

Let $T > 0$. Suppose we are given an $\mathcal{F}_T$ -measurable random variable ξ , a progressively measurable function $f : \Omega \times [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ and two adapted càdlàg processes L, U such that $L_t \leq U_t$, $t \in [0, T]$. Roughly speaking, by a solution of the reflected BSDE with terminal condition ξ , generator f and barriers L, U we mean a quadruple (Y, K, A, M) of càdlàg adapted processes such that Y is of Doob's class D, K, A are increasing processes such that $K_0 = A_0 = 0$, M is a local martingale with $M_0 = 0$, and a.s. we have

$$\begin{cases} Y_t = \xi + \int_t^T f(s, Y_s) ds + \int_t^T dK_s - \int_t^T dA_s - \int_t^T dM_s, & t \in [0, T] \\ L_t \leq Y_t \leq U_t, & t \in [0, T], \\ \int_0^T (Y_{t-} - L_{t-}) dK_t = \int_0^T (U_{t-} - Y_{t-}) dA_t = 0. \end{cases} \quad (1.1)$$

In most papers devoted to reflected BSDEs with two barriers it is assumed that L, U satisfy one of the following conditions:

- (a) between U and L one can find a process X such that X is a difference of nonnegative càdlàg supermartingales (so-called Mokobodzki condition), or
- (b) $L_t < U_t$ and $L_{t-} < U_{t-}$ for $t \in [0, T]$ (i.e. the barriers are completely separated).

Problem (1.1) under assumption (a) is studied thoroughly in Klimisak [8]. Among other things, in [8] it is proved that if f is continuous and monotone with respect to y and satisfies mild integrability conditions (see hypotheses (H1)–(H4) in Section 2), then there exists a unique solution of (1.1).

A drawback to assumption (a), and one of the main reason why more explicit condition (b) is considered, is that (a) may be sometimes difficult to check. Unfortunately, equations with barriers satisfying (b) are more difficult to deal with. At present, all the existing results on equations with barriers satisfying (b) concern the case where the underlying filtration is Brownian (see Hamadène and Hassani [5], Hamadène, Hassani and Ouknine [6]) or is generated by a Brownian motion and an independent Poisson random measure (see Hamadène and Wang [7]). Moreover, in [5, 6, 7] it is assumed that f is Lipschitz continuous and the data (including barriers) are L^2 -integrable. Recently, in [4], in the case of Brownian filtration, an existence and uniqueness result was proved for equations with separated continuous barriers, L^1 data and Lipschitz continuous generator.

Our main theorem says that under the assumptions on ξ, f from [8] and càdlàg barriers U, L satisfying (b) and such that L^+, U^- are of class D there exists a unique solution of (1.1). Thus we extend the results from [8] to barriers satisfying (b) and at the same time we generalize the results of [5, 6, 7, 4] to equations with general filtration and less regular data. It is worth pointing out that as a simple corollary to our existence result (it suffices to consider the generator $f \equiv 0$) one gets the following result from the general theory of stochastic processes: if two càdlàg processes L, U are completely separated and L^+, U^- are of class D, then there exists a semimartingale of class D between U and L .

The main idea of the proof of our main result is to reduce the problem with completely separated barriers to the problem with barriers satisfying the Mokobodzki condition, and then apply the results of [8]. Such a reduction is possible locally (we use here some modification of a construction from [3]) and enables us to obtain solutions of (1.1) on stochastic intervals of the form $[0, \tau_n]$, where $\{\tau_n\}$ is some stationary sequence of stopping times. These local solutions can be put together to get the solution of (1.1) on $[0, T]$. The last step involves some technicalities, but in general our proof is short and rather simple. In our opinion, it is much simpler than the proof for equations with the underlying Brownian-Poisson filtration and L^2 data given in [7].

The paper is organized as follows. In Section 2 we review some results from [8] concerning reflected BSDEs with one barrier. The proof of the main result is given in Section 3. Finally, in Section 4 we give an application of results of Section 3 to zero-sum Dynkin game with payoff function determined by ξ, f and L, U .

Notation. Let $T > 0$, and let $(\Omega, \mathcal{F}, \mathbb{F} = \{\mathcal{F}_t\}_{t \in [0, T]}, P)$ be a filtered probability space with filtration satisfying the usual assumptions of completeness and right continuity. By $\mathcal{T}$ we denote the set of all $\mathbb{F}$ -stopping times such that $\tau \leq T$, and by $\mathcal{T}_t, t \in [0, T]$, the set of $\tau \in \mathcal{T}$ such that $P(\tau \geq t) = 1$.

By $\mathcal{V}$ we denote the set of all $\mathbb{F}$ -progressively measurable processes of finite variation, and by $\mathcal{V}^1$ the subset of $\mathcal{V}$ consisting of all processes V such that $E|V|_T < \infty$, where $|V|_T$ stands for the variation of V on $[0, T]$. $\mathcal{V}_0$ is the subset of $\mathcal{V}$ consisting of all processes V such that $V_0 = 0$, $\mathcal{V}_0^+$ (resp. ${}^p\mathcal{V}_0^+$) is the subset of $\mathcal{V}_0$ of all increasing processes (resp. predictable increasing processes). $\mathcal{M}$ (resp. $\mathcal{M}_{loc}$) denotes the set of all $\mathbb{F}$ -martingales (resp. local martingales). By $L^1(\mathbb{F})$ we denote the space of all $\mathbb{F}$ -progressively measurable

processes X such that $E \int_0^T |X_t| dt < \infty$, and by $L^1(\mathcal{F}_T)$ the space of all $\mathcal{F}_T$ -measurable random variables ξ such that $E|\xi| < \infty$.

For a stochastic process X we set $X^+ = X \vee 0$, $X^- = -(X \wedge 0)$ and $X_{t-} = \lim_{s \nearrow t} X_s$ with the convention that $X_{0-} = X_0$. We also adopt the convention that $\int_a^b = \int_{(a,b]}$.

2 BSDEs with one reflecting barrier

In what follows ξ is an $\mathcal{F}_T$ -measurable random variable, L, U are $\mathbb{F}$ -progressively measurable càdlàg processes, $V \in \mathcal{V}_0$ and $f : \Omega \times [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ is a measurable function such that $f(\cdot, y)$ is an $\mathbb{F}$ -progressively measurable process for every $y \in \mathbb{R}$ (for brevity, in our notation we omit the dependence of f on $\omega \in \Omega$).

We will need the following assumptions on ξ and f :

(H1) There exists a constant $\mu \in \mathbb{R}$ such that for almost every $t \in [0, T]$ and all $y, y' \in \mathbb{R}$,

$$(f(t, y) - f(t, y'))(y - y') \leq \mu |y - y'|^2,$$

(H2) $[0, T] \ni t \mapsto f(t, y) \in L^1(0, T)$ for every $y \in \mathbb{R}$,

(H3) the function $\mathbb{R} \ni y \mapsto f(t, y)$ is continuous for almost every $t \in [0, T]$,

(H4) $\xi \in L^1(\mathcal{F}_T)$, $V \in \mathcal{V}_0 \cap \mathcal{V}^1$, $f(\cdot, 0) \in L^1(\mathcal{F})$.

Recall that a stochastic process X on $[0, T]$ is said to be of class D if $\{X_\tau : \tau \in \mathcal{T}\}$ is a uniformly integrable family of random variables.

Definition 2.1. We say that a triple (Y, K, M) of càdlàg processes is a solution of the reflected BSDE with terminal condition ξ , generator $f + dV$ and lower barrier L (RBSDE $(\xi, f + dV, L)$ for short) if

(a) Y is a process of class D, $K \in {}^p\mathcal{V}_0^+$, $M \in \mathcal{M}_{loc}$ with $M_0 = 0$,

(b) $L_t \leq Y_t$, $t \in [0, T]$ P -a.s.,

(c) $\int_0^T (Y_{t-} - L_{t-}) dK_t = 0$,

(d) $Y_t = \xi + \int_t^T f(s, Y_s) ds + \int_t^T dV_s + \int_t^T dK_s - \int_t^T dM_s$, $t \in [0, T]$ P -a.s.

Definition 2.2. We say that a triple (Y, K, M) of càdlàg processes is a solution of the reflected BSDE with terminal condition ξ , generator $f + dV$ and upper barrier U (RBSDE $(\xi, f + dV, U)$ for short) if

(a) Y is a process of class D, $A \in {}^p\mathcal{V}_0^+$, $M \in \mathcal{M}_{loc}$ with $M_0 = 0$,

(b) $Y_t \leq U_t$ for $t \in [0, T]$ P -a.s.,

(c) $\int_0^T (U_{t-} - Y_{t-}) dA_t = 0$,

(d) $Y_t = \xi + \int_t^T f(s, Y_s) ds + \int_t^T dV_s - \int_t^T dA_s - \int_t^T dM_s$, $t \in [0, T]$ P -a.s.

Our motivations for considering reflected equations involving a finite variation process V comes from the theory of partial differential equations with measure data. In these applications V is an additive functional of a Markov process in the Revuz correspondence with some smooth measure see [9, 10, 11]

In the theorem below we recall some results on reflecting BSDEs with one barrier proved in [8]. They will play important role in the proof of our main result in Section 3.

Theorem 2.3. Assume that L^+, U^- are of class D and (H1)–(H4) are satisfied.

- (i) There exists a unique solution $(\underline{Y}, \underline{K}, \underline{M})$ of $\underline{RBSDE}(\xi, f + dV, L)$. Moreover, if $(\underline{Y}^n, \underline{M}^n)$, $n \in \mathbb{N}$, are solutions of BSDEs of the form

$$\underline{Y}_t^n = \xi + \int_t^T f(s, \underline{Y}_s^n) ds + \int_t^T dV_s + \int_t^T n(L_s - \underline{Y}_s^n)^+ ds - \int_t^T d\underline{M}_s^n,$$

then $\underline{Y}_t^n \nearrow \underline{Y}_t$, $t \in [0, T]$ P -a.s.

- (ii) There exists a unique solution $(\tilde{Y}, \tilde{A}, \tilde{M})$ of $\overline{RBSDE}(\xi, f + dV, U)$. Moreover, if $(\tilde{Y}^n, \tilde{M}^n)$, $n \in \mathbb{N}$, are solutions of BSDEs of the form

$$\tilde{Y}_t^n = \xi + \int_t^T f(s, \tilde{Y}_s^n) ds + \int_t^T dV_s - \int_t^T n(\tilde{Y}_s^n - U_s)^+ ds - \int_t^T d\tilde{M}_s^n,$$

then $\tilde{Y}_t^n \nearrow \tilde{Y}_t$, $t \in [0, T]$ P -a.s.

PROOF. Part (i) is proved in [8, Theorem 4.1] under the assumption that L is of class D. The following argument shows that in fact it suffices to assume that L^+ is of class D. Let (Y^0, M^0) be a solution of the BSDE

$$Y_t^0 = \xi + \int_t^T f(s, Y_s^0) ds + \int_t^T dV_s - \int_t^T dM_s^0, \quad t \in [0, T], \quad (2.1)$$

and let $L^\varepsilon = L \vee (Y^0 - \varepsilon)$ for some $\varepsilon > 0$. If L^+ is of class D then L^ε is of class D, because Y^0 is of class D. Therefore by [8, Theorem 4.1] there exists a solution $(Y^\varepsilon, K^\varepsilon, M^\varepsilon)$ of $\underline{RBSDE}(\xi, f + dV, L^\varepsilon)$ such that $K^\varepsilon \in \mathcal{V}_0^+$. In particular,

$$Y_t^\varepsilon = \xi + \int_t^T f(s, Y_s^\varepsilon) ds + \int_t^T dV_s + \int_t^T dK_s^\varepsilon - \int_t^T dM_s^\varepsilon, \quad t \in [0, T] \quad (2.2)$$

and

$$Y^\varepsilon \geq L^\varepsilon \geq L. \quad (2.3)$$

By (2.1), (2.2) and [9, Proposition 2.1], $Y^\varepsilon \geq Y^0$. Hence $\mathbb{1}_{\{Y_{t-}^\varepsilon > L_{t-}\}} = \mathbb{1}_{\{Y_{t-}^\varepsilon > L_{t-}^\varepsilon\}}$, $t \in [0, T]$, and consequently

$$\begin{aligned} \int_0^T (Y_{t-}^\varepsilon - L_{t-}) dK_t^\varepsilon &= \int_0^T (Y_{t-}^\varepsilon - L_{t-}) \mathbb{1}_{\{Y_{t-}^\varepsilon > L_{t-}\}}(t) dK_t^\varepsilon \\ &= \int_0^T (Y_{t-}^\varepsilon - L_{t-}) \mathbb{1}_{\{Y_{t-}^\varepsilon > L_{t-}^\varepsilon\}}(t) dK_t^\varepsilon = 0, \end{aligned} \quad (2.4)$$

the last equality being a consequence of the fact that $\int_0^T \mathbb{1}_{\{Y_{t-}^\varepsilon > L_{t-}^\varepsilon\}}(t) dK_t^\varepsilon = 0$. By (2.2)–(2.4) the triple $(\underline{Y}, \underline{K}, \underline{M}) = (Y^\varepsilon, K^\varepsilon, M^\varepsilon)$ is a solution of $\underline{RBSDE}(\xi, f + dV, L)$. Uniqueness follows from [8, Corollary 2.2.]. This proves the first part of (i). Observe now that the first component of the solution of $\underline{RBSDE}(\xi, 0, L)$ is a supermartingale of class D majorizing L . Therefore to prove that $\underline{Y}_t^n \nearrow \underline{Y}_t$, $t \in [0, T]$, it suffices to repeat step by step the proof of [8, Theorem 4.1]. Since the proof of (ii) is analogous to that of (i), we omit it. $\square$

3 BSDEs with two reflecting barriers

In this section ξ, f, V and U, L are as in Section 2. We also assume that $L_t \leq U_t$ for $t \in [0, T]$ P -a.s.

Definition 3.1. We say that a quadruple (Y, K, A, M) of càdlàg processes is a solution of the reflected BSDE with terminal condition ξ , generator $f + dV$, lower barrier L and upper barrier U ($(Y, K, A, M) = RBSDE(\xi, f + dV, L, U)$ for short) if

(LU1) Y is a process of class D, $A, K \in \mathcal{V}_0^+$, $M \in \mathcal{M}_{loc}$ with $M_0 = 0$,

(LU2) $L_t \leq Y_t \leq U_t$ for $t \in [0, T]$ P -a.s.,

(LU3) $\int_0^T (Y_{t-} - L_{t-}) dK_t = \int_0^T (U_{t-} - Y_{t-}) dA_t = 0$,

(LU4) $Y_t = \xi + \int_t^T f(s, Y_s) ds + \int_t^T dV_s + \int_t^T d(K_s - A_s) - \int_t^T dM_s$, $t \in [0, T]$ P -a.s.

Let us consider following conditions for barriers L, U .

(B1) $L_t < U_t$ and $L_{t-} < U_{t-}$ for $t \in [0, T]$.

(B2) L^+, U^- are processes of class D.

A sequence $\{\tau_n\} \subset \mathcal{T}$ is called of stationary type, if

$$P\left(\liminf_{n \rightarrow \infty} \{\tau_n = T\}\right) = 1.$$

The following lemma is an extension of [3, Remark 3.4].

Lemma 3.2. Assume that L, U are of class D and satisfy (B1). Then there exists a process $H \in \mathcal{V}$ such that $L_t \leq H_t \leq U_t$, $t \in [0, T]$ P -a.s. Moreover, there exists a sequence $\{\tau_n\} \subset \mathcal{T}$ of stationary type such that $E|H|_{\tau_n} < \infty$ for every $n \in \mathbb{N}$.

PROOF. Let $\tau_0 = 0$, and for $n \in \mathbb{N}$ set

$$\tau_n = \inf \left\{ t > \tau_{n-1} : \frac{L_{\tau_{n-1}} + U_{\tau_{n-1}}}{2} > U_t \quad \text{or} \quad \frac{L_{\tau_{n-1}} + U_{\tau_{n-1}}}{2} < L_t \right\} \wedge T.$$

Obviously $\{\tau_n\}$ is nondecreasing. We shall show that it is increasing up to T . To see this, let us first observe that

$$P(\tau_n = \tau_{n+1} < T) = 0, \quad n \in \mathbb{N} \cup \{0\}. \quad (3.1)$$

Indeed, suppose that $\omega \in \{\tau_n = \tau_{n+1} < T\}$. Then there exists a sequence $\{t_m\}$ such that $t_m \searrow \tau_n(\omega)$ and for every $m \in \mathbb{N}$,

$$\frac{L_{\tau_n(\omega)}(\omega) + U_{\tau_n(\omega)}(\omega)}{2} > U_{t_m}(\omega) \quad \text{or} \quad \frac{L_{\tau_n(\omega)}(\omega) + U_{\tau_n(\omega)}(\omega)}{2} < L_{t_m}(\omega).$$

Since L and U are right-continuous, this implies that

$$\frac{L_{\tau_n(\omega)}(\omega) + U_{\tau_n(\omega)}(\omega)}{2} \geq U_{\tau_n(\omega)}(\omega) \quad \text{or} \quad \frac{L_{\tau_n(\omega)}(\omega) + U_{\tau_n(\omega)}(\omega)}{2} \leq L_{\tau_n(\omega)}(\omega).$$

Hence $L_{\tau_n(\omega)}(\omega) = U_{\tau_n(\omega)}(\omega)$. Since the barriers satisfy (B1), this shows (3.1). We can now prove that $\{\tau_n\}$ is of stationary type. Suppose, for contradiction, that there is $\tau \in \mathcal{T}$ such that $\tau_n \nearrow \tau$ and $P(\bigcap_{n=1}^{\infty} \{\tau_n < \tau\}) > 0$. Then

$$P\left(\frac{L_{\tau_{n-1}} + U_{\tau_{n-1}}}{2} \rightarrow \frac{L_{\tau-} + U_{\tau-}}{2}\right) > 0.$$

This implies that $P(L_{\tau-} \geq \frac{L_{\tau-} + U_{\tau-}}{2} \geq U_{\tau-}) > 0$, hence that $P(L_{\tau-} \geq U_{\tau-}) > 0$, in contradiction with (B1). Thus $\{\tau_n\}$ is of stationary type. Set

$$H_t = \sum_{n=1}^{\infty} \frac{L_{\tau_{n-1}} + U_{\tau_{n-1}}}{2} \mathbb{1}_{[\tau_{n-1}, \tau_n)}(t), \quad t \in [0, T].$$

Then $L_t \leq H_t \leq U_t$, $t \in [0, T]$ P -a.e., and $H \in \mathcal{V}$, because $\{\tau_n\}$ is of stationary type. Moreover, for each $n \in \mathbb{N}$,

$$E|H|_{\tau_n} = \sum_{k=1}^n E \left| \frac{U_{\tau_k} + L_{\tau_k}}{2} - \frac{U_{\tau_{k-1}} + L_{\tau_{k-1}}}{2} \right|,$$

which is finite because L, U are of class D. $\square$

The following example shows that in general there is no H between barriers such that $E|H|_T$ is finite.

Example 3.3. Let $\{B_n\}$ be a partition of Ω such that $P(B_n) = Cn^{-2}$ with $C = 6\pi^{-2}$, and let $h : [0, 1) \rightarrow \mathbb{R}$ be defined by the following formula

$$h_t = \begin{cases} \frac{1}{2}, & t \in [1 - \frac{1}{2n+1}, 1 - \frac{1}{2n+2}), \quad n \in \mathbb{N} \cup \{0\}, \\ -\frac{3}{2}, & t \in [1 - \frac{1}{2n}, 1 - \frac{1}{2n+1}), \quad n \in \mathbb{N}. \end{cases}$$

Set $T = 1$ and

$$L_t = \sum_{n=1}^{\infty} h_{t \wedge (1 - \frac{1}{n+1})} \mathbb{1}_{B_n}, \quad U_t = L_t + 1, \quad t \in [0, T].$$

One can check that L, U satisfy the assumptions of Lemma 3.2. Therefore there exists a process $H \in \mathcal{V}$ such that $L_t \leq H_t \leq U_t$, $t \in [0, T]$ P -a.s.

Consider now an arbitrary process $\bar{H} \in \mathcal{V}$ such that $L_t \leq \bar{H}_t \leq U_t$, $t \in [0, T]$ P -a.s. By the construction of the barriers L and U ,

$$|\bar{H}|_T \mathbb{1}_{B_n} \geq \sum_{t \in [0, T]} (|U_t - L_{t-}| \wedge |U_{t-} - L_t|) \mathbb{1}_{\{L_t - L_{t-} \neq 0\}}(t) \mathbb{1}_{B_n} = n \mathbb{1}_{B_n}.$$

Hence

$$E|\bar{H}|_T = \sum_{n=1}^{\infty} E|\bar{H}|_T \mathbb{1}_{B_n} \geq \sum_{n=1}^{\infty} n P(B_n) = \sum_{n=1}^{\infty} \frac{C}{n} = \infty.$$

Before proving our main result, we first introduce some additional notation. Assume that ξ, f satisfy (H1)–(H4), and L, U are of class D and satisfy (B1). Set

$$\underline{f}_m(t, y) = f(t, y) - m(y - U_t)^+, \quad \bar{f}_n(t, y) = f(t, y) + n(L_t - y)^+.$$

Then $\underline{f}_m, \bar{f}_n$ also satisfy (H1)–(H4), because $y \mapsto n(L_t - y)^+$ and $y \mapsto m(y - U_t)^+$ are Lipschitz continuous for $t \in [0, T]$ and L, U are of class D. By Theorem 2.3, for each $n \in \mathbb{N}$ there exists a unique solution $(\bar{Y}^n, \bar{A}^n, \bar{M}^n)$ of $\underline{RBSDE}(\xi, \bar{f}_n + dV, U)$, and for each $m \in \mathbb{N}$ there exists a unique solution $(\underline{Y}^m, \underline{K}^m, \underline{M}^m)$ of $\underline{RBSDE}(\xi, \underline{f}_m + dV, L)$. Therefore

$$\begin{aligned} \bar{Y}_t^n &= \xi + \int_t^T f(s, \bar{Y}_s^n) ds + \int_t^T dV_s \\ &\quad + \int_t^T n(L_s - \bar{Y}_s^n)^+ ds - \int_t^T d\bar{A}_s^n - \int_t^T d\bar{M}_s^n \leq U_t \end{aligned} \quad (3.2)$$

and

$$\begin{aligned}\underline{Y}_t^m &= \xi + \int_t^T f(s, \underline{Y}_s^m) ds + \int_t^T dV_s \\ &\quad - \int_t^T m(\underline{Y}_s^m - U_s)^+ ds + \int_t^T d\underline{K}_s^m - \int_t^T d\underline{M}_s^m \geq L_t\end{aligned}$$

for $t \in [0, T]$. The function $(t, y) \mapsto f(t, y) - m(y - U_t)^+ + n(L_s - y)^+$ also satisfies (H1)–(H4), so by [8, Theorem 2.7], for any $n, m \in \mathbb{N}$ there exists a solution $(Y^{n,m}, M^{n,m})$ of the BSDE

$$\begin{aligned}Y_t^{n,m} &= \xi + \int_t^T f(s, Y_s^{n,m}) ds + \int_t^T dV_s + \int_t^T n(L_s - Y_s^{n,m})^+ ds \\ &\quad - \int_t^T m(Y_s^{n,m} - U_s)^+ ds - \int_t^T dM_s^{n,m}, \quad t \in [0, T].\end{aligned}$$

By Theorem 2.3, for each $m \in \mathbb{N}$ the sequence $\{Y^{n,m}\}_n$ is nondecreasing, for each $n \in \mathbb{N}$ the sequence $\{Y^{n,m}\}_m$ is nonincreasing, and

$$\underline{Y}_t^m = \sup_{n \in \mathbb{N}} Y_t^{n,m} = \lim_{n \rightarrow \infty} Y_t^{n,m}, \quad \overline{Y}_t^n = \inf_{m \in \mathbb{N}} Y_t^{n,m} = \lim_{m \rightarrow \infty} Y_t^{n,m}, \quad t \in [0, T]$$

In particular, for all $n, m \in \mathbb{N}$ we have

$$\overline{Y}_t^n \leq Y_t^{n,m} \leq \underline{Y}_t^m, \quad t \in [0, T]. \quad (3.3)$$

By [8, Proposition 2.1] the sequence $\{\overline{Y}^n\}$ is nondecreasing, whereas the sequence $\{\underline{Y}^m\}$ is nonincreasing. Set

$$\underline{Y}_t = \inf_{m \in \mathbb{N}} \underline{Y}_t^m = \lim_{m \rightarrow \infty} \underline{Y}_t^m, \quad \overline{Y}_t = \sup_{n \in \mathbb{N}} \overline{Y}_t^n = \lim_{n \rightarrow \infty} \overline{Y}_t^n. \quad (3.4)$$

Since $\underline{Y}^m \geq L$ for all $m \in \mathbb{N}$ and $\overline{Y}^n \leq U$ for all $n \in \mathbb{N}$, we have

$$\underline{Y}_t \geq L_t, \quad \overline{Y}_t \leq U_t, \quad t \in [0, T] \quad P\text{-a.s.}$$

Also note that from (3.3) and (3.4) and monotonicity of the sequences $\{\overline{Y}^n\}$, $\{\underline{Y}^m\}$ it follows that

$$\overline{Y}^0 \leq \overline{Y}^n \leq \overline{Y} \leq \underline{Y} \leq \underline{Y}^m \leq \underline{Y}^0. \quad (3.5)$$

Since $\overline{Y}^0$ and $\underline{Y}^0$ are solution of reflected BSDEs, they are processes of class D.

Lemma 3.4. *Assume (H1), (H2). Then for every $r > 0$*

$$t \mapsto \sup_{|y| \leq r} f(t, y) \in L^1(0, T).$$

PROOF. By (H1), for all $y \in [-r, r]$, $t \in [0, T]$ we have

$$f(t, y) \geq f(t, r) - 2\mu r, \quad f(t, y) \leq f(t, -r) + 2\mu r.$$

Hence

$$\sup_{|y| \leq r} |f(t, y)| \leq |f(t, -r) + 2\mu r| \vee |f(t, r) - 2\mu r|.$$

It suffices to use (H2) to complete the proof. $\square$

Lemma 3.5. *Let (Y, K, A, M) be a solution of $RBSDE(\xi, f + dV, L, U)$, and let $\tau \in \mathcal{T}$. If $\xi \in L^1(\mathcal{F}_\tau)$, $f(t, y)\mathbb{1}_{(\tau, T]}(t) = 0$ for all $y \in \mathbb{R}$ and $t \in [0, T]$, and*

$$V_t = V_{t \wedge \tau}, \quad L_t = L_{t \wedge \tau}, \quad U_t = U_{t \wedge \tau}, \quad t \in [0, T], \quad (3.6)$$

then

$$K_t = K_{t \wedge \tau}, \quad A_t = A_{t \wedge \tau}, \quad M_t = M_{t \wedge \tau}, \quad t \in [0, T]. \quad (3.7)$$

PROOF. By (LU4),

$$Y_{t \wedge \tau} - Y_t = \int_{t \wedge \tau}^t dK_s - \int_{t \wedge \tau}^t dA_s - \int_{t \wedge \tau}^t dM_s. \quad (3.8)$$

Let $\{\zeta_n\}$ be a fundamental sequence for the local martingale M , and let $\sigma \in \mathcal{T}$. Applying the Tanaka–Meyer formula we get

$$\begin{aligned} (Y_{(\sigma \wedge \zeta_n) \wedge \tau} - Y_{\sigma \wedge \zeta_n})^+ &\leq \int_{(\sigma \wedge \zeta_n) \wedge \tau}^{\sigma \wedge \zeta_n} \mathbb{1}_{\{Y_{(s \wedge \tau)-} > Y_{s-}\}} dK_s \\ &\quad - \int_{(\sigma \wedge \zeta_n) \wedge \tau}^{\sigma \wedge \zeta_n} \mathbb{1}_{\{Y_{(s \wedge \tau)-} > Y_{s-}\}} dA_s - \int_{(\sigma \wedge \zeta_n) \wedge \tau}^{\sigma \wedge \zeta_n} \mathbb{1}_{\{Y_{(s \wedge \tau)-} > Y_{s-}\}} dM_s \\ &\leq \int_{(\sigma \wedge \zeta_n) \wedge \tau}^{\sigma \wedge \zeta_n} \mathbb{1}_{\{Y_\tau > Y_{s-}\}} dK_s - \int_{(\sigma \wedge \zeta_n) \wedge \tau}^{\sigma \wedge \zeta_n} \mathbb{1}_{\{Y_\tau > Y_{s-}\}} dM_s. \end{aligned}$$

Taking the expectation and then letting $n \rightarrow \infty$ yields

$$E(Y_{\sigma \wedge \tau} - Y_\sigma)^+ \leq E \int_{\sigma \wedge \tau}^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} dK_s.$$

On the other hand, by (LU3) and (3.6),

$$\int_{\sigma \wedge \tau}^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} dK_s = \int_{\sigma \wedge \tau}^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_{s-}\}} dK_s = \int_{\sigma \wedge \tau}^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s.$$

Hence

$$E(Y_{\sigma \wedge \tau} - Y_\sigma)^+ \leq E \int_{\sigma \wedge \tau}^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s. \quad (3.9)$$

From now on we consider the stopping time σ defined by

$$\sigma = \inf\{t > \tau : Y_{t \wedge \tau} > Y_t\} \wedge T.$$

Observe that

$$Y_{t \wedge \tau} \mathbb{1}_{\{t < \sigma\}} \leq Y_t \mathbb{1}_{\{t < \sigma\}}, \quad t \in [0, T]. \quad (3.10)$$

Set

$$B_T = \left\{ \int_\tau^T \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s > 0 \right\}.$$

Since $\mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} \leq \mathbb{1}_{\{L_\tau < Y_\tau\}}$, we have

$$B_T \subset \{L_\tau < Y_\tau\}. \quad (3.11)$$

From (LU3), (3.10), (3.11) and the fact that $L_t = L_{t \wedge \tau}$ for $t \in [0, T]$ it follows that

$$\mathbb{1}_{B_T} \cdot \int_{\tau \wedge \sigma}^\sigma dK_s = 0. \quad (3.12)$$

By (3.9) and (3.12),

$$\begin{aligned} E(Y_\tau - Y_\sigma)^+ &\leq E \int_\tau^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s \\ &= E \left(\mathbb{1}_{B_T} \int_\tau^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s \right) = 0. \end{aligned}$$

By the above,

$$E((Y_\tau - Y_\sigma)^+ \mathbb{1}_{\{\sigma = T\}}) = 0, \quad E((Y_\tau - Y_\sigma)^+ \mathbb{1}_{\{\sigma < T\}}) = 0. \quad (3.13)$$

Suppose that $P(\sigma = T) = 1$. Then by (3.10) and the first equality in (3.13), $(Y_{t \wedge \tau} - Y_t)^+ = 0$ P -a.s. for $t \in [0, T]$. We now prove that

$$P(\sigma < T) = 0. \quad (3.14)$$

By the second equality in (3.13),

$$P(\{Y_\tau \leq Y_\sigma\} \cap \{\sigma < T\}) = P(\sigma < T). \quad (3.15)$$

Observe that from the definition of σ and the fact that $L_t = L_{t \wedge \tau}$ for $t \in [0, T]$ it follows that

$$\{\sigma < T\} \subset \{L_\tau < Y_\tau\}. \quad (3.16)$$

Set

$$\zeta = \inf \left\{ t > \sigma : Y_t < \frac{Y_\tau + L_\tau}{2} \right\}.$$

By right-continuity of Y and (3.16), $Y_\zeta \mathbb{1}_{\{\sigma < T\}} \leq \frac{Y_\tau + L_\tau}{2} \mathbb{1}_{\{\sigma < T\}}$. By this and (3.16),

$$P(\{Y_\zeta < Y_\tau\} \cap \{\sigma < T\}) = P(\sigma < T). \quad (3.17)$$

Furthermore, from (3.11), the definition of ζ and (LU3) it follows that

$$0 \leq \mathbb{1}_{B_T} \cdot \int_\sigma^\zeta dK_s \leq \mathbb{1}_{\{L_\tau < Y_\tau\}} \cdot \int_\sigma^\zeta dK_s = \mathbb{1}_{\{L_\tau < Y_\tau\}} \mathbb{1}_{\{\sigma < \zeta\}} \cdot \int_\sigma^\zeta dK_s = 0. \quad (3.18)$$

Observe that by the definition of the set B_T ,

$$E \left(\int_\tau^\zeta \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s \right) = E \left(\mathbb{1}_{B_T} \int_\tau^\zeta \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s \right).$$

By the above equality, (3.12) and (3.18),

$$\begin{aligned} E \left(\int_\tau^\zeta \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s \right) &= E \left(\mathbb{1}_{B_T} \int_\tau^\sigma \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s \right) \\ &\quad + E \left(\mathbb{1}_{B_T} \int_\sigma^\zeta \mathbb{1}_{\{Y_\tau > Y_{s-}\}} \mathbb{1}_{\{Y_{s-} = L_\tau\}} dK_s \right) = 0. \end{aligned}$$

This when combined with (3.9) with σ replaced by ζ gives $E(Y_\tau - Y_\zeta)^+ = 0$. Consequently, $E(Y_\tau - Y_\zeta)^+ \mathbb{1}_{\{\sigma < T\}} = 0$, which when combined with (3.17) proves (3.14). Thus $(Y_{t \wedge \tau} - Y_t)^+ = 0$, $t \in [0, T]$ P -a.s. Applying the Tanaka–Meyer formula to the process $(Y_{t \wedge \tau} - Y_t)^-$ and using similar arguments one can prove that $(Y_{t \wedge \tau} - Y_t)^- = 0$, $t \in [0, T]$ P -a.s. Hence

$$Y_t = Y_{t \wedge \tau}, \quad t \in [0, T]. \quad (3.19)$$

From (3.8) and (3.19) we obtain

$$0 = \int_{t \wedge \tau}^t dK_s - \int_{t \wedge \tau}^t dA_s - \int_{t \wedge \tau}^t dM_s,$$

which implies (3.7). $\square$

Theorem 3.6. Assume that (H1)–(H4), (B1), (B2) are satisfied. Then there exists a unique solution (Y, K, A, M) of $RBSDE(\xi, f + dV, L, U)$. Moreover, $Y = \overline{Y} = \underline{Y}$.

PROOF. By [8, Corollary 3.2] there exists at most one solution of $RBSDE(\xi, f + dV, L, U)$, so it suffices to prove the existence of a solution. To this end, we first assume additionally that L, U are of class D. Then by Lemma 3.2 there exists $H \in \mathcal{V}$ such that $L_t \leq H_t \leq U_t$, $t \in [0, T]$ P -a.s. and $H_{\cdot \wedge \tau'_k} \in \mathcal{V}^1$ for some sequence $\{\tau'_k\}$ of stationary type. Set

$$\tau_k = \tau'_k \wedge \delta_k \quad (3.20)$$

and $H^{(k)} = H_{\cdot \wedge \tau_k}$, where

$$\delta_k = \inf \left\{ t \geq 0 : \int_0^t f^-(s, H_s) ds > k \right\} \wedge T.$$

Observe that $H^{(k)} \in \mathcal{V}^1$ and by Lemma 3.4 $\{\tau_k\}$ is of stationary type. The rest of the proof we divide into 5 steps.

Step 1. We show the existence of a solution of $RBSDE(\xi, f + dV, L, U)$ on stochastic intervals $[0, \tau_k]$. Set

$$\begin{aligned} U^{(k)} &= U_{\cdot \wedge \tau_k}, & L^{(k)} &= L \mathbb{1}_{[0, \tau_k)} + (L_{\tau_k} \wedge \overline{Y}_{\tau_k}) \mathbb{1}_{[\tau_k, T]}, \\ \xi^{(k)} &= \overline{Y}_{\tau_k}, & f^{(k)}(\cdot, y) &= f(\cdot, y) \mathbb{1}_{[0, \tau_k]}, & V^{(k)} &= V_{\cdot \wedge \tau_k}, \end{aligned}$$

where $\overline{Y}$ is defined by (3.4). By (3.5) $\xi^{(k)} \in L^1(\mathcal{F}_T)$. Also observe that $L_T^{(k)} \leq \xi^{(k)} \leq U_T^{(k)}$ and $L_t^{(k)} \leq H_t^{(k)} \leq U_t^{(k)}$, $t \in [0, T]$. Therefore by [8, Theorem 3.3] there exists a unique solution $(Y^{(k)}, K^{(k)}, A^{(k)}, M^{(k)})$ of $RBSDE(\xi^{(k)}, f^{(k)} + dV^{(k)}, L^{(k)}, U^{(k)})$ such that

$$EK_T^{(k)} < \infty, \quad EA_T^{(k)} < \infty. \quad (3.21)$$

In particular, we have

$$\begin{aligned} Y_t^{(k)} &= \xi^{(k)} + \int_t^T f^{(k)}(s, Y_s^{(k)}) ds + \int_t^T dV_s^{(k)} \\ &\quad + \int_t^T dK_s^{(k)} - \int_t^T dA_s^{(k)} - \int_t^T dM_s^{(k)} \end{aligned} \quad (3.22)$$

for $t \in [0, T]$. By Lemma 3.5,

$$(Y_t^{(k)}, K_t^{(k)}, A_t^{(k)}, M_t^{(k)}) = (Y_{t \wedge \tau_k}^{(k)}, K_{t \wedge \tau_k}^{(k)}, A_{t \wedge \tau_k}^{(k)}, M_{t \wedge \tau_k}^{(k)}), \quad t \in [0, T]. \quad (3.23)$$

Step 2. We are going to show for every $\tau \in \mathcal{T}$,

$$Y_\tau^{(k)} = \overline{Y}_{\tau \wedge \tau_k}. \quad (3.24)$$

By Theorem 2.3, for each $n \in \mathbb{N}$ there exists a unique solution $(Y^{(k),n}, A^{(k),n}, M^{(k),n})$ of $\overline{RBSDE}(\xi^{(k)}, f^{(k),n} + dV^{(k)}, U^{(k)})$ with $f^{(k),n}(t, y) = f^{(k)}(t, y) + n(L_t^{(k)} - y)^+$ and the triple $(Y^{(k),n}, A^{(k),n}, M^{(k),n})$ satisfies the equation

$$\begin{aligned} Y_t^{(k),n} &= \xi^{(k)} + \int_t^T f^{(k)}(s, Y_s^{(k),n}) ds + \int_t^T dV_s^{(k)} \\ &\quad + \int_t^T n(L_s^{(k)} - Y_s^{(k),n})^+ ds - \int_t^T dA_s^{(k),n} - \int_t^T dM_s^{(k),n}. \end{aligned} \quad (3.25)$$

and by [8, Theorem 3.3]

$$Y^{(k),n} \nearrow Y^{(k)}. \quad (3.26)$$

To this end, we write $\tilde{Y}_t^n = Y_t^{(k),n} - \bar{Y}_t^n$, $\tilde{A}_t^n = A_t^{(k),n} - \bar{A}_t^n$, $\tilde{M}_t^n = M_t^{(k),n} - \bar{M}_t^n$. By (3.2), (3.25) and the Tanaka-Meyer formula, for all $\zeta, \tau \in \mathcal{T}$ we have

$$\begin{aligned} \tilde{Y}_{\tau \wedge \zeta \wedge \tau_k}^{n,+} &\leq \tilde{Y}_{\zeta \wedge \tau_k}^{n,+} + \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} (f^{(k)}(s, Y_s^{(k),n}) - f^{(k)}(s, \bar{Y}_s^n)) ds \\ &\quad + \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} n \left((L_s^{(k)} - Y_s^{(k),n})^+ - (L_s - \bar{Y}_s^n)^+ \right) ds \\ &\quad - \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} d\tilde{A}_s^n - \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^{n,+} > 0\}} d\tilde{M}_s^n. \end{aligned}$$

By this and (H1),

$$\begin{aligned} \tilde{Y}_{\tau \wedge \zeta \wedge \tau_k}^{n,+} &\leq \tilde{Y}_{\zeta \wedge \tau_k}^{n,+} + \mu \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \tilde{Y}_s^{n,+} ds \\ &\quad + \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} n \left((L_s^{(k)} - Y_s^{(k),n})^+ - (L_s - \bar{Y}_s^n)^+ \right) ds \\ &\quad + \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} d\bar{A}_s^n - \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} d\bar{M}_s^n. \end{aligned} \quad (3.27)$$

Since $y \mapsto (L_s - y)^+$ is nonincreasing and $L^{(k)} \mathbb{1}_{[0, \tau_k)} = L \mathbb{1}_{[0, \tau_k)}$, we have

$$\int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} ((L_s^{(k)} - Y_s^{(k),n})^+ - (L_s - \bar{Y}_s^n)^+) ds \leq 0. \quad (3.28)$$

Since $Y_{t \wedge \tau_k}^{(k),n} \leq U_{t \wedge \tau_k}^{(k)} = U_t^{(k)}$ and $\bar{Y}_{t \wedge \tau_k}^n \leq U_{t \wedge \tau_k} = U_t$, we have $\bar{Y}_{t \wedge \tau_k}^n \leq Y_{t \wedge \tau_k}^{(k),n} \vee \bar{Y}_{t \wedge \tau_k}^n \leq U_t^{(k)}$. Hence

$$\begin{aligned} \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^{n,+} > 0\}} d\bar{A}_s^n &\leq \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} \frac{Y_{s-}^{(k),n} \vee \bar{Y}_{s-}^n - \bar{Y}_{s-}^n}{\tilde{Y}_{s-}^n} d\bar{A}_s^n \\ &\leq \liminf_{m \rightarrow \infty} m \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > \frac{1}{m}\}} (U_{s-} - \bar{Y}_{s-}^n) d\bar{A}_s^n = 0. \end{aligned} \quad (3.29)$$

By (3.27)–(3.29),

$$\tilde{Y}_{\tau \wedge \zeta \wedge \tau_k}^{n,+} \leq \tilde{Y}_{\zeta \wedge \tau_k}^{n,+} + \mu \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \tilde{Y}_s^{n,+} ds - \int_{\tau \wedge \zeta \wedge \tau_k}^{\zeta \wedge \tau_k} \mathbb{1}_{\{\tilde{Y}_{s-}^n > 0\}} d\tilde{M}_s^n$$

for any $\tau, \zeta \in \mathcal{T}$. Let $\{\zeta_m\}$ be a fundamental sequence for the local martingale $\tilde{M}^n$. Replacing ζ by ζ_m in the above inequality and then taking the expectation we obtain

$$E\tilde{Y}_{\tau \wedge \zeta_m \wedge \tau_k}^{n,+} \leq E\tilde{Y}_{\zeta_m \wedge \tau_k}^{n,+} + \mu E \int_{\tau \wedge \zeta_m \wedge \tau_k}^{\zeta_m \wedge \tau_k} \tilde{Y}_s^{n,+} ds.$$

The processes $Y^{(k)}, \bar{Y}^n$ are of class D as solutions of reflected BSDEs. Consequently, $\tilde{Y}^{n,+}$ is of class D. Therefore letting $m \rightarrow \infty$ in the above inequality we get

$$E\tilde{Y}_{\tau \wedge \tau_k}^{n,+} \leq E\tilde{Y}_{\tau_k}^{n,+} + \mu E \int_{\tau \wedge \tau_k}^{\tau_k} \tilde{Y}_s^{n,+} ds \quad (3.30)$$

for all $\tau \in \mathcal{T}$. Observe that

$$\int_{(\tau \vee t) \wedge \tau_k}^{\tau_k} \tilde{Y}_s^{n,+} ds = \int_{((\tau \wedge \tau_k) \vee t) \wedge \tau_k}^{\tau_k} \tilde{Y}_s^{n,+} ds = \int_t^T \tilde{Y}_s^{n,+} \mathbb{1}_{[\tau \wedge \tau_k, \tau_k]}(s) ds \leq \int_t^T \tilde{Y}_s^{n,+} ds.$$

From the above inequality and (3.30) with τ replaced by $\tau \vee t$ it follows that

$$E\tilde{Y}_{(\tau \vee t) \wedge \tau_k}^{n,+} \leq E\tilde{Y}_{\tau_k}^{n,+} + \mu \int_t^T E\tilde{Y}_{(\tau \vee s) \wedge \tau_k}^{n,+} ds, \quad \tau \in \mathcal{T}, t \in [0, T].$$

Applying Gronwall's inequality to the mapping $t \mapsto E\tilde{Y}_{(\tau \vee t) \wedge \tau_k}^{n,+}$ gives

$$E\tilde{Y}_{(\tau \vee t) \wedge \tau_k}^{n,+} \leq e^{\mu T} E\tilde{Y}_{\tau_k}^{n,+} \leq e^{\mu T} E|Y_{\tau_k}^{(k),n} - \bar{Y}_{\tau_k}^n|, \quad t \in [0, T]. \quad (3.31)$$

By (3.4), $\bar{Y}_{\tau_k}^n \nearrow \bar{Y}_{\tau_k} = \xi^{(k)}$, whereas by (3.26) and (3.23), $Y_{\tau_k}^{(k),n} \nearrow Y_{\tau_k}^{(k)} = \xi^{(k)}$. Hence, by the monotone convergence theorem,

$$E|Y_{\tau_k}^{(k),n} - \xi^{(k)}| \rightarrow 0, \quad E|\bar{Y}_{\tau_k}^n - \xi^{(k)}| \rightarrow 0.$$

Therefore applying Fatou's lemma and then (3.31) with $t = T$ we obtain

$$E \liminf_{n \rightarrow \infty} \tilde{Y}_{(\tau \wedge \tau_k)}^{n,+} \leq \liminf_{n \rightarrow \infty} E\tilde{Y}_{(\tau \wedge \tau_k)}^{n,+} \leq \liminf_{n \rightarrow \infty} e^{\mu T} (E|Y_{\tau_k}^{(k),n} - \xi^{(k)}| + E|\bar{Y}_{\tau_k}^n - \xi^{(k)}|) = 0.$$

But $\tilde{Y}_{\tau \wedge \tau_k}^n \rightarrow Y_{\tau \wedge \tau_k}^{(k)} - \bar{Y}_{\tau \wedge \tau_k} = Y_{\tau}^{(k)} - \bar{Y}_{\tau \wedge \tau_k}$. Hence $E(Y_{\tau}^{(k)} - \bar{Y}_{\tau \wedge \tau_k})^+ = 0$. In much the same way one can show that $E(Y_{\tau}^{(k)} - \bar{Y}_{\tau \wedge \tau_k})^- = 0$, which completes the proof of (3.24). By (3.24) and the optional cross-section theorem [2, p. 138-IV, (86) Theorem] the processes $Y^{(k)}$ and $\bar{Y}_{\cdot \wedge \tau_k}$ are indistinguishable. In particular, $\bar{Y}_{\cdot \wedge \tau_k}$ has càdlàg trajectories. By the same method we show that $Y^{(k)}$ and $\underline{Y}_{\cdot \wedge \tau_k}$ are indistinguishable.

Step 3. In this step we define a solution on $[0, T]$. By Step 2, for every $k \in \mathbb{N}$,

$$Y_{t \wedge \tau_k}^{(k)} = \bar{Y}_{t \wedge \tau_k} = \bar{Y}_{t \wedge \tau_k \wedge \tau_{k+1}} = Y_{t \wedge \tau_k}^{(k+1)}, \quad t \in [0, T]. \quad (3.32)$$

By (3.22), (3.32) and uniqueness of the semimartingale decomposition,

$$(Y_{t \wedge \tau_k}^{(k+1)}, K_{t \wedge \tau_k}^{(k+1)}, A_{t \wedge \tau_k}^{(k+1)}, M_{t \wedge \tau_k}^{(k+1)}) = (Y_{t \wedge \tau_k}^{(k)}, K_{t \wedge \tau_k}^{(k)}, A_{t \wedge \tau_k}^{(k)}, M_{t \wedge \tau_k}^{(k)}), \quad t \in [0, T].$$

Therefore we may define processes Y, K, A, M on $[0, T]$ by

$$Y_t = Y_t^{(k)}, \quad K_t = K_t^{(k)}, \quad A_t = A_t^{(k)}, \quad M_t = M_t^{(k)}, \quad t \in [0, \tau_k]. \quad (3.33)$$

By Step 2, $Y_{\tau \wedge \tau_k} = \underline{Y}_{\tau \wedge \tau_k} = \bar{Y}_{\tau \wedge \tau_k}$ for all $\tau \in \mathcal{T}$ and $k \in \mathbb{N}$, so letting $k \rightarrow \infty$ gives $Y_{\tau} = \bar{Y}_{\tau}$ for $\tau \in \mathcal{T}$. Hence, by the cross-section theorem,

$$Y = \bar{Y}.$$

The quadruple (Y, K, A, M) is a solution of $RBSDE(\xi, f + dV, L, U)$. Indeed, from (3.22), (3.33) and stationarity of $\{\tau_k\}$ it follows that (Y, K, A, M) satisfies (LU1) and (LU4). Moreover, from the fact that $(Y^{(k)}, K^{(k)}, A^{(k)}, M^{(k)})$ is a solution of $RBSDE(\xi^{(k)}, f^{(k)} + dV^{(k)}, L^{(k)}, U^{(k)})$ and (3.33) it follows that $L_{t \wedge \tau_k} \leq Y_{t \wedge \tau_k} \leq U_{t \wedge \tau_k}$ for $t \in [0, T]$ P -a.s. and

$$\int_0^{\tau_k} (Y_{t-} - L_{t-}) dK_t = \int_0^{\tau_k} (U_{t-} - Y_{t-}) dA_t = 0$$

for $k \in \mathbb{N}$. Since $\{\tau_k\}$ is of stationary type, this implies (LU2) and (LU3).

Step 4. Repeating arguments from step 3 and step 4 for $\xi^{(k)} = \underline{Y}_{\tau_k}$ we prove $\underline{Y} = Y$, where (Y, K, A, M) is a solution of $RBSDE(\xi, f + dV, L, U)$. Therefore, by the uniqueness of solution, $Y = \overline{Y} = \underline{Y}$.

Step 5. We now show how to dispense with the assumption that L, U are of class D.

Let $\underline{Y}, \tilde{Y}$ be processes appearing in Theorem 2.3. By [9, Proposition 2.1],

$$\tilde{Y}_t \leq \underline{Y}_t, \quad t \in [0, T], \quad P\text{-a.s.}$$

Let $\varepsilon > 0$, and let $L_t^\varepsilon = L_t \vee (\tilde{Y}_t - \varepsilon)$, $U_t^\varepsilon = U_t \wedge (\underline{Y}_t + \varepsilon)$. If L, U satisfy (B1), (B2), then also $L^\varepsilon, U^\varepsilon$ satisfy (B1) and are processes of class (D). By steps 1-3 there exists unique solution (Y, K, A, M) of $RBSDE(\xi, f + dV, L^\varepsilon, U^\varepsilon)$. Analogously to the proof of Theorem 2.3 it could be verified, that (Y, K, A, M) is also solution of $RBSDE(\xi, f + dV, L, U)$. $\square$

Corollary 3.7. *Assume that L, U satisfy (B1), (B2). Then there exists a semimartingale Y of class D such that $L_t \leq Y_t \leq U_t$, $t \in [0, T]$ P -a.s.*

PROOF. It is enough to consider $\xi = L_T^+ \wedge U_T$, $f \equiv 0$, $V \equiv 0$, and apply Theorem 3.6. $\square$

Remark 3.8. Let $\{\tau_n\}$ be a sequence defined by (3.20). If there exists $k_0 \in \mathbb{N}$ such that

$$P(\tau_{k_0} = T) = 1, \quad (3.34)$$

then by Step 3 of the proof of Theorem 3.6, $(Y, K, A, M) = (Y^{(k_0)}, K^{(k_0)}, A^{(k_0)}, M^{(k_0)})$ is a solution of $RBSDE(\xi, f + dV, L, U)$. Furthermore, by (3.21), $EK_T < \infty$ and $EA_T < \infty$, and by [9, Lemma 2.3], $f(\cdot, Y) \in L^1(\mathcal{F})$. Also note that a sufficient condition for (3.34) to hold is the following: there is $H \in \mathcal{V}^1$ such that $L_t \leq H_t \leq U_t$, $t \in [0, T]$, and $t \rightarrow f(t, H_t)$ is bounded from below.

The following example shows that in general EK_T and EA_T need not be finite even if $f \equiv 0$ and $V \equiv 0$.

Example 3.9. Let $\mathbb{F}$ be a Brownian filtration, and let L, U be defined as in Example 3.3. Set $\xi = (L_T + U_T)/2$ and $f \equiv 0$, $V \equiv 0$. By Theorem 3.6 there exists a unique solution (Y, K, A, M) of $RBSDE(\xi, 0, L, U)$. In particular,

$$Y_t = \xi + \int_t^T dK_s - \int_t^T dA_s - \int_t^T dM_s, \quad t \in [0, T].$$

Let $\tau_n = 1 - \frac{1}{n}$. Since the filtration is Brownian, $\Delta M_{\tau_n} = 0$ P -a.s. for every $n \in \mathbb{N}$. Hence

$$\Delta Y_{\tau_n} = \Delta A_{\tau_n} - \Delta K_{\tau_n}, \quad n \in \mathbb{N}.$$

In fact, by (LU2), (LU3) and the definitions of L and U , $\Delta Y_{\tau_m} = \Delta A_{\tau_m}$ if m is even and $\Delta Y_{\tau_m} = -\Delta K_{\tau_m}$ if m is odd. From this and the fact that $L \leq Y \leq U$ it follows that

$$P(\{\Delta A_{\tau_m} \geq 1\} \cap B_n) = Cn^{-2}, \quad 2 \leq m \leq n+1$$

when m is even, and

$$P(\{\Delta K_{\tau_m} \geq 1\} \cap B_n) = Cn^{-2}, \quad 2 \leq m \leq n+1$$

when m is odd. Hence

$$EK_T = E|K|_T = \sum_{n=1}^{\infty} E|K|_T \mathbb{1}_{B_n} \geq \sum_{n=2}^{\infty} \frac{n-1}{2} P(B_n) = C \sum_{n=2}^{\infty} \frac{n-1}{2n^2} = \infty$$

and

$$EA_T = E|A|_T = \sum_{n=1}^{\infty} E|A|_T \mathbb{1}_{B_n} \geq \sum_{n=2}^{\infty} \frac{n-1}{2} P(B_n) = C \sum_{n=2}^{\infty} \frac{n-1}{2n^2} = \infty.$$

4 Dynkin games

In this section we consider a certain stochastic game of stopping called Dynkin game. For an interpretation of notions which we define below (payoff function, lower and upper value of the game) we defer the reader to [1].

Let L, U be càdlàg processes of class D such that $L_t \leq U_t$, $t \in [0, T]$ P -a.s., and let f, ξ, V be as in Section 3. Also assume that conditions (H1)–(H4) are satisfied. Consider a stopping game with payoff function

$$R_t(\sigma, \tau) = \int_t^{\sigma \wedge \tau} f(s, Y_s) ds + \int_t^{\sigma \wedge \tau} dV_s + \xi \mathbb{1}_{\{\sigma \wedge \tau = T\}} + L_\tau \mathbb{1}_{\{\tau < T, \tau \leq \sigma\}} + U_\sigma \mathbb{1}_{\{\sigma < \tau\}}, \quad \sigma, \tau \in \mathcal{T}_t, \quad (4.1)$$

where (Y, K, A, M) is a solution of $RBSDE(\xi, f + dV, L, U)$ such that $K, A \in \mathcal{V}_0^1$. By Remark 3.8 such solution exists if (B1), (B2) and (3.34) are satisfied.

The lower value $\underline{V}$ and the upper value $\overline{V}$ of the stochastic game corresponding to R are defined by

$$\underline{V}_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \operatorname{ess\,inf}_{\sigma \in \mathcal{T}_t} E(R_t(\sigma, \tau) | \mathcal{F}_t), \quad \overline{V}_t = \operatorname{ess\,inf}_{\sigma \in \mathcal{T}_t} \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} E(R_t(\sigma, \tau) | \mathcal{F}_t).$$

We say that the game has a value if $\underline{V}_t = \overline{V}_t$, $t \in [0, T]$ P -a.s.

Lemma 4.1. *Let $\{\tau_n\}$ be a sequence of stopping times such that $\tau_n \nearrow \tau$ P -a.s., and*

$$P(\liminf_{n \rightarrow \infty} \{\tau_n = \tau\}) = 1. \quad (4.2)$$

Then for every $\sigma \in \mathcal{T}_t$, $E(R_t(\sigma, \tau_n) | \mathcal{F}_t) \rightarrow E(R_t(\sigma, \tau) | \mathcal{F}_t)$ P -a.s. as $n \rightarrow \infty$.

PROOF. By (4.1) and (4.2), $R_t(\sigma, \tau_n) \rightarrow R_t(\sigma, \tau)$ P -a.s. Since V, L, U are of class D and $E|\xi| + E \int_0^T |f(t, Y_t)| dt < \infty$, we conclude from (4.1) that $\{R_t(\sigma, \tau_n)\}_{n \in \mathbb{N}}$ is a uniformly integrable family of random variables. Therefore the desired convergence follows from [13, Theorem 1.3]. $\square$

Theorem 4.2. *Let the assumptions of Theorem 3.6 hold, and moreover (3.34) is satisfied. Then the stochastic game corresponding to the payoff function (4.1) has the value equal to the first component of the solution of $RBSDE(\xi, f + dV, L, U)$, i.e.*

$$Y_t = \underline{V}_t = \overline{V}_t, \quad t \in [0, T] \quad P\text{-a.s.} \quad (4.3)$$

PROOF. By [12, Lemma 5.3], to show that the game has a value it suffices to prove that for any $\varepsilon > 0$ and $t \in [0, T]$ there exist $\sigma_t^\varepsilon, \tau_t^\varepsilon \in \mathcal{T}_t$ such that for all $\sigma, \tau \in \mathcal{T}_t$,

$$-\varepsilon + E(R_t(\sigma_t^\varepsilon, \tau) | \mathcal{F}_t) \leq E(R_t(\sigma, \tau_t^\varepsilon) | \mathcal{F}_t) + \varepsilon. \quad (4.4)$$

To show (4.4), we set $\sigma_t^\varepsilon = \inf\{s > t : Y_s \geq U_s - \varepsilon\} \wedge T$. Observe that $Y_{s-} < U_{s-}$ for $t < s \leq \sigma_t^\varepsilon$, and hence, by (LU3),

$$A_s \mathbb{1}_{(t, \sigma_t^\varepsilon]}(s) = A_{\sigma_t^\varepsilon} \mathbb{1}_{(t, \sigma_t^\varepsilon]}(s), \quad s \in [0, T]. \quad (4.5)$$

Clearly, for any $\tau \in \mathcal{T}_t$,

$$\{\sigma_t^\varepsilon = T\} \subset \{\tau \leq \sigma_t^\varepsilon\}, \quad \{\tau > \sigma_t^\varepsilon\} \subset \{\sigma_t^\varepsilon < T\}.$$

It follows that on the set $\{\tau \leq \sigma_t^\varepsilon\}$, by (4.5), we have

$$\begin{aligned} R_t(\sigma_t^\varepsilon, \tau) &= \int_t^\tau f(s, Y_s) ds + \int_t^\tau dV_s + \xi \mathbb{1}_{\{\tau=T\}} + L_\tau \mathbb{1}_{\{\tau < T\}} \\ &\leq \int_t^\tau f(s, Y_s) ds + \int_t^\tau dV_s + \xi \mathbb{1}_{\{\tau=T\}} + Y_\tau \mathbb{1}_{\{\tau < T\}} + \int_t^\tau dK_s - \int_t^\tau dA_s \\ &\leq Y_t + \int_t^\tau dM_s, \end{aligned}$$

whereas on $\{\tau > \sigma_t^\varepsilon\}$ we have

$$\begin{aligned} R_t(\sigma_t^\varepsilon, \tau) &= \int_t^{\sigma_t^\varepsilon} f(s, Y_s) ds + \int_t^{\sigma_t^\varepsilon} dV_s + U_{\sigma_t^\varepsilon} \mathbb{1}_{\{\sigma_t^\varepsilon < \tau\}} \\ &\leq \int_t^{\sigma_t^\varepsilon} f(s, Y_s) ds + \int_t^{\sigma_t^\varepsilon} dV_s + Y_{\sigma_t^\varepsilon} + \int_t^{\sigma_t^\varepsilon} dK_s - \int_t^{\sigma_t^\varepsilon} dA_s + \varepsilon \\ &= Y_t + \int_t^{\sigma_t^\varepsilon} dM_s + \varepsilon. \end{aligned}$$

Hence

$$R_t(\sigma_t^\varepsilon, \tau) = R_t(\sigma_t^\varepsilon, \tau) \mathbb{1}_{\{\tau \leq \sigma_t^\varepsilon\}} + R_t(\sigma_t^\varepsilon, \tau) \mathbb{1}_{\{\tau > \sigma_t^\varepsilon\}} \leq Y_t + \int_t^{\sigma_t^\varepsilon \wedge \tau} dM_s + \varepsilon$$

P -a.s. Let $\{\zeta_n\}$ be a fundamental sequence for the local martingale M , and let $\tau_n = \tau \wedge \zeta_n$. Then $\{\tau_n\}$ satisfies the assumptions of Lemma 4.1 and

$$E(R_t(\sigma_t^\varepsilon, \tau \wedge \zeta_n) | \mathcal{F}_t) \leq E\left(Y_t + \int_t^{\sigma_t^\varepsilon \wedge \tau \wedge \zeta_n} dM_s + \varepsilon | \mathcal{F}_t\right) = Y_t + \varepsilon.$$

Letting $n \rightarrow \infty$ and using Lemma 4.1 we obtain

$$E(R_t(\sigma_t^\varepsilon, \tau) | \mathcal{F}_t) \leq Y_t + \varepsilon. \quad (4.6)$$

Now, let us consider the stopping time $\tau_t^\varepsilon = \inf\{s > t : Y_s \leq L_s + \varepsilon\} \wedge T$. Analysis similar to that in the proof of (4.6) shows that for any $\varepsilon > 0$ and $\sigma \in \mathcal{T}_t$,

$$E(R_t(\sigma, \tau_t^\varepsilon) | \mathcal{F}_t) \geq Y_t - \varepsilon. \quad (4.7)$$

Combining (4.6) with (4.7) we see that for any $\varepsilon > 0$,

$$-\varepsilon + E(R_t(\sigma_t^\varepsilon, \tau) | \mathcal{F}_t) \leq Y_t \leq E(R_t(\sigma, \tau_t^\varepsilon) | \mathcal{F}_t) + \varepsilon. \quad (4.8)$$

Thus (4.4) is satisfied, and, in consequence, the game has a value. Moreover, from (4.8) and the definitions of $\underline{V}, \overline{V}$ it follows that $-\varepsilon + \overline{V}_t \leq Y_t \leq \underline{V}_t + \varepsilon$, $t \in [0, T]$, for $\varepsilon > 0$. Since we already know that the game has a value, this implies (4.3). $\square$

Note that Dynkin games were studied, in different contexts, by several authors. For results related to Theorem 4.2 we defer the reader to [1, 8, 12, 14, 15] and references given there.

References

- [1] J. Cvitanic, I. Karatzas, *Backward stochastic differential equations with reflection and Dynkin games*, Ann. Probab. 24 (1996), 2024–2056.
- [2] C. Dellacherie, P.-A. Meyer, *Probabilities and potential*, North-Holland, Paris 1978.
- [3] A. Falkowski, *Stochastic differential equations with respect to processes of finite p-variation* (in Polish), PhD thesis, Nicolaus Copernicus University, 2015.
- [4] I. Hassairi, *Existence and uniqueness for D-solutions of reflected BSDEs with two barriers without Mokobodzki's condition*, Commun. Pure Appl. Anal. 15 (2016), 1139–1156.
- [5] S. Hamadène, M. Hassani, *BSDEs with two reflecting barriers: the general result*, Probab. Theory Relat. Fields 132 (2005), 237–264.
- [6] S. Hamadène, M. Hassani, Y. Ouknine, *Backward SDEs with two rcll reflecting barriers without Mokobodzki's hypothesis*, Bull. Sci. Math. 134 (2010), 874–899.
- [7] S. Hamadène, H. Wang, *BSDEs with two RCLL reflecting obstacles driven by Brownian motion and Poisson measure and a related mixed zero-sum game*, Stochastic Process. Appl. 119 (2009), 2881–2912.
- [8] T. Klimsiak, *Reflected BSDEs on filtered probability spaces*, Stochastic Process. Appl. 125 (2015), 4204–4241.
- [9] T. Klimsiak, A. Rozkosz, *Dirichlet forms and semilinear elliptic equations with measure data*, J. Funct. Anal. 265 (2013), 890–925.
- [10] T. Klimsiak, A. Rozkosz, *Obstacle problem for semilinear parabolic equations with measure data*, J. Evol. Equ. 15 (2015), 457–491.
- [11] T. Klimsiak, A. Rozkosz, *Semilinear elliptic equations with measure data and quasi-regular Dirichlet forms*, Colloq. Math. 145 (2016), 35–67.
- [12] J.-P. Lepeltier, M. Xu, *Reflected backward stochastic differential equations with two RCLL barriers*, ESAIM: Probab. Stat. 11 (2007), 3–22.
- [13] R. Liptzer, A. N. Shiryaev, *Statistics of Random Processes*, Springer-Verlag, New York 2001.
- [14] Ł. Stettner, *On a general zero-sum stochastic game with optimal stopping*, Probab. Math. Statist. 3 (1982), 103–112.
- [15] J. Zabczyk, *Stopping games for symmetric Markov process*, Probab. Math. Statist. 4 (1984), 185–196.